

	2022					
	Number	Number, low CI	Number, high CI	Percent	Percent, low CI	Percent, high CI
Fibromyalgia						
Total, by sex	571,183	520,135	622,231	1.7%	1.6%	1.9%
Males	93,846	73,151	114,541	0.6%	0.4%	0.7%
Females	477,337	430,707	523,967	2.9%	2.6%	3.2%
Total, by age	571,183	520,135	622,231	1.7%	1.6%	1.9%
Less than 65	370,506	326,295	414,718	1.4%	1.3%	1.6%
65 or older	200,677	176,275	225,078	2.9%	2.5%	3.2%
Chronic fatigue syndrome						
Total, by sex	443,896	396,240	491,553	1.3%	1.2%	1.5%
Males	158,173	128,607	187,738	1.0%	0.8%	1.2%
Females	285,723	248,672	322,774	1.7%	1.5%	1.9%
Total, by age	443,896	396,240	491,553	1.3%	1.2%	1.5%
Less than 65	300,561	260,431	340,691	1.2%	1.0%	1.3%
65 or older	143,335	119,565	167,106	2.0%	1.7%	2.4%
Both conditions						
Total, by sex	150,630	124,572	32,499	0.5%	0.4%	0.5%
Males	22,458	12,418	32,499	0.1%	0.1%	0.2%
Females	128,172	104,037	152,306	0.8%	0.6%	0.9%
Total, by age	150,630	124,572	32,499	0.5%	0.4%	0.5%
Less than 65	97,177	76916.94683	117,436	0.4%	0.3%	0.5%
65 or older	53,453	37,160	69,747	0.8%	0.5%	1.0%
Multiple Chemical Sensitivities						
Total, by sex	563,018	508,870	617,166	1.7%	1.5%	1.9%
Males	182,775	149,920	215,630	1.1%	0.9%	1.3%
Females	380,243	334,899	425,587	2.3%	2.0%	2.6%
Total, by age	563,018	508,870	617,166	1.7%	1.5%	1.9%
Less than 65	395,056	345,344	444,767	1.5%	1.3%	1.7%
65 or older	167,962	146,131	189,794	2.4%	2.1%	1.7%

	2021					
	Number	Number, low CI	Number, high CI	Percent	Percent, low CI	Percent, high CI
Fibromyalgia						
Total, by sex	526,396	470,440	582,351	1.6%	1.4%	1.8%
Males	106,110	71,420	140,801	0.7%	0.4%	0.9%
Females	420,285	376,402	464,168	2.6%	2.3%	2.8%
Total, by age	526,396	470,440	582,351	1.6%	1.4%	1.8%
Less than 65	333,010	281,883	384,137	1.3%	1.1%	1.5%
65 or older	193,386	169,526	217,245	2.9%	2.5%	3.2%
Chronic fatigue syndrome						
Total, by sex	426,775	377,810	475,740	1.3%	1.2%	1.5%
Males	156,620	127,482	185,758	1.0%	0.8%	1.2%
Females	270,154	229,575	310,734	1.6%	1.4%	1.9%
Total, by age	426,775	377,810	475,740	1.3%	1.2%	1.5%
Less than 65	299,368	253,852	344,883	1.2%	1.0%	1.3%
65 or older	127,407	107,483	147,331	1.9%	1.6%	2.2%
Both conditions						
Total, by sex	94,182	75,751	112,612	0.3%	0.2%	0.3%
Males	11,963	4,337	19,589	0.1%	0.0%	0.1%
Females	82,219	65,596	98,842	0.5%	0.4%	0.6%
Total, by age	94,182	75,751	112,612	0.3%	0.2%	0.3%
Less than 65	64,369	47,886	80,853	0.3%	0.2%	0.3%
65 or older	29,813	21,184	38,441	0.4%	0.3%	0.6%
Multiple Chemical Sensitivities						
Total, by sex	954,401	877,212	1,031,591	2.9%	2.7%	3.2%
Males	318,266	260,528	376,005	2.0%	1.6%	2.3%
Females	636,135	581,459	690,811	3.9%	3.5%	4.2%
Total, by age	954,401	877,212	1,031,591	2.9%	2.7%	3.2%
Less than 65	654,109	580,903	727,315	2.5%	2.3%	2.8%
65 or older	318,266	269,577	331,008	4.5%	4.0%	4.9%

	2019					
	Number	Number, low CI	Number, high CI	Percent	Percent, low CI	Percent, high CI
Fibromyalgia						
Total, by sex	572,200	521,500	622,800	1.8%	1.6%	2.0%
Males	114,800	88,000	141,600	0.7%	0.6%	0.9%
Females	457,400	415,900	498,800	2.8%	2.6%	3.1%
Total, by age	572,200	521,500	622,800	1.8%	1.6%	2.0%
Less than 65	377,300	331,900	422,700	1.5%	1.3%	1.7%
65 or older	194,800	171,300	218,400	3.1%	2.7%	3.4%
Chronic fatigue syndrome						
Total, by sex	462,700	418,500	507,000	1.5%	1.3%	1.6%
Males	153,000	126,800	179,100	1.0%	0.8%	1.1%
Females	309,800	274,400	345,200	1.9%	1.7%	2.1%
Total, by age	462,700	418,500	507,000	1.5%	1.3%	1.6%
Less than 65	325,400	286,100	364,700	1.3%	1.1%	1.4%
65 or older	137,300	117,600	157,000	2.2%	1.9%	2.5%
Both conditions						
Total, by sex	132,100	110,200	154,100	0.4%	0.3%	0.5%
Males	18,300	10,600	26,000	0.1%	0.1%	0.2%
Females	113,800	93,600	134,100	0.7%	0.6%	0.8%
Total, by age	132,100	110,200	154,100	0.4%	0.3%	0.5%
Less than 65	88,900	70,300	107,500	0.4%	0.3%	0.4%
65 or older	43,200	31,900	54,600	0.7%	0.5%	0.9%

	2016					
	Number	Number, low CI	Number, high CI	Percent	Percent, low CI	Percent, high CI
Fibromyalgia						
Total, by sex	522,000	473,700	570,300	1.7%	1.5%	1.9%
Males	106,300	83,500	129,100	0.7%	0.6%	0.9%
Females	415,700	372,900	458,500	2.7%	2.4%	3.0%
Total, by age	522,000	473,700	570,300	1.7%	1.5%	1.9%
Less than 65	317,500	279,000	356,100	1.3%	1.1%	1.4%
65 or older	204,500	174,600	234,300	3.6%	3.1%	4.1%
Chronic fatigue syndrome						
Total, by sex	581,600	525,300	637,900	1.9%	1.7%	2.1%
Males	208,300	175,000	241,600	1.4%	1.2%	1.6%
Females	373,400	326,300	420,400	2.4%	2.1%	2.7%
Total, by age	581,600	525,300	637,900	1.9%	1.7%	2.1%
Less than 65	434,700	383,200	486,300	1.7%	1.5%	2.0%
65 or older	146,900	124,100	169,600	2.6%	2.2%	3.0%
Both conditions						
Total, by sex	139,000	114,400	163,600	0.5%	0.4%	0.5%
Males	20,200	11,500	28,900	0.1%	0.1%	0.2%
Females	118,900	96,500	141,200	0.8%	0.6%	0.9%
Total, by age	139,000	114,400	163,600	0.5%	0.4%	0.5%
Less than 65	93,600	74,700	112,500	0.4%	0.3%	0.5%
65 or older	45,400	30,300	60,600	0.8%	0.5%	1.1%

	2015					
	Number	Number, low CI	Number, high CI	Percent	Percent, low CI	Percent, high CI
Fibromyalgia						
Total, by sex	493,600	445,200	542,000	1.6%	1.5%	1.8%
Males	97,200	69,500	125,000	0.7%	0.5%	0.8%
Females	396,400	357,800	434,900	2.6%	2.3%	2.8%
Total, by age	493,600	445,200	542,000	1.6%	1.5%	1.8%
Less than 65	335,800	294,800	376,900	1.4%	1.2%	1.5%
65 or older	157,800	133,900	181,600	2.9%	2.4%	3.3%
Chronic fatigue syndrome						
Total, by sex	561,500	506,800	616,100	1.9%	1.7%	2.0%
Males	195,100	160,900	229,200	1.3%	1.1%	1.5%
Females	366,400	322,000	410,800	2.4%	2.1%	2.7%
Total, by age	561,500	506,800	616,100	1.9%	1.7%	2.0%
Less than 65	404,500	356,400	452,600	1.6%	1.4%	1.8%
65 or older	156,900	132,500	181,400	2.9%	2.4%	3.3%
Both conditions						
Total, by sex	138,400	107,700	169,200	0.5%	0.4%	0.6%
Males	23,400	1,700	45,000	0.2%	0.0%	0.3%
Females	115,100	92,900	137,200	0.8%	0.6%	0.9%
Total, by age	138,400	107,700	169,200	0.5%	0.4%	0.6%
Less than 65	103,700	76,600	130,800	0.4%	0.3%	0.5%
65 or older	34,700	22,000	47,400	0.6%	0.4%	0.9%