



	A	B	C	D	E	F	G	H	I
1									
2									
3									
4									
5		2020							
6		Number		Number, low CI	Number, high CI	Percent		Percent, low CI	Percent, high CI
7	Fibromyalgia								
8	Total, by sex	526,400		470,400	582,400	1.6%		1.4%	1.8%
9	Males	106,100	E	71,400	140,800	0.7%	E	0.4%	0.9%
10	Females	420,300		376,400	464,200	2.6%		2.3%	2.8%
11	Total, by age	526,400		470,400	582,400	1.6%		1.4%	1.8%
12	Less than 65	333,000		281,900	384,100	1.3%		1.1%	1.5%
13	65 or older	193,400		169,500	217,200	2.9%		2.5%	3.2%
14	Chronic fatigue syndrome								
15	Total, by sex	426,800		377,800	475,700	1.3%		1.2%	1.5%
16	Males	156,600		127,500	185,800	1.0%		0.8%	1.2%
17	Females	270,200		229,600	310,700	1.6%		1.4%	1.9%
18	Total, by age	426,800		377,800	475,700	1.3%		1.2%	1.5%
19	Less than 65	299,400		253,900	344,900	1.2%		1.0%	1.3%
20	65 or older	127,400		107,500	147,300	1.9%		1.6%	2.2%
21	Both conditions								
22	Total, by sex	94,200		75,800	112,600	0.3%		0.2%	0.3%
23	Males	12,000	E	4,300	19,600	0.1%	E	0.0%	0.1%
24	Females	82,200		65,600	98,800	0.5%		0.4%	0.6%
25	Total, by age	94,200		75,800	112,600	0.3%		0.2%	0.3%
26	Less than 65	64,400		47,900	80,900	0.3%		0.2%	0.3%
27	65 or older	29,800		21,200	38,400	0.4%		0.3%	0.6%
28	Multiple Chemical Sensitivities								
29	Total, by sex	954,400		877,200	1,031,600	2.9%		2.7%	3.2%
30	Males	318,300		260,500	376,000	2.0%		1.6%	2.3%
31	Females	636,100		581,500	690,800	3.9%		3.5%	4.2%
32	Total, by age	954,400		877,200	1,031,600	2.9%		2.7%	3.2%
33	Less than 65	654,100		580,900	727,300	2.5%		2.3%	2.8%
34	65 or older	300,300		269,600	331,000	4.5%		4.0%	4.9%